

Honour Chemistry: Practice Final Exam (Semester 2)

Part A: Multiple Choice and Numerical Response

1. Which of the following equations is associated with the largest energy change per mole of fluorine?
- A. $F_2(g) \rightarrow F_2(l)$
B. ${}^{19}_9F + {}^4_2He \rightarrow {}^1_0n + {}^{22}_{11}Na$
C. $2 F_2(g) + 2 H_2O(l) \rightarrow O_2(g) + 4 HF(aq)$
D. $CH_4(g) + 2 Cl_2(g) + 2 F_2(g) \rightarrow CCl_2F_2(g) + 2 HCl(g) + 2 HF(g)$
2. Which of the following molecular properties is a main component of the chemical potential energy of matter?
- A. Vibrational motion
B. Intramolecular bonding
C. Movement from place to place
D. Rotation about the molecules' centre of mass
3. When one mole of sodium bicarbonate is formed from its elements, 947.7 kJ of heat energy is released into the surroundings. This enthalpy change can be represented as
- A. $Na(s) + \frac{1}{2} H_2(g) + C(s) + \frac{3}{2} O_2(g) \rightarrow NaHCO_3(s) + 947.7 \text{ kJ}$
B. $Na(s) + \frac{1}{2} H_2(g) + C(s) + \frac{3}{2} O_2(g) + 947.7 \text{ kJ} \rightarrow NaHCO_3(s)$
C. $Na^+(aq) + HCO_3^-(aq) \rightarrow NaHCO_3(s) + 947.7 \text{ kJ}$
D. $Na^+(aq) + HCO_3^-(aq) + 947.7 \text{ kJ} \rightarrow NaHCO_3(s)$

Use the following information to answer the next question.

Cold packs are commonly used by athletes to reduce swelling caused by injury. The packs consist of two plastic pouches: an inner pouch that contains a chemical and an outer pouch that contains water. When the inner pouch is broken, the chemical and water mix, which causes the pack to feel cold.

Statements

- 1 Ice is considerably less expensive than are commercial cold packs.
- 2 Ammonium nitrate is commonly used in cold packs because its heat of solution is endothermic.
- 3 The disposal of cold packs poses a landfill concern.
- 4 Durability and flexibility are design requirements for the plastic outer pouch.



Numerical Response


1. The statements above that reflect an ecological, scientific, economic and technological perspective are, respectively, _____, _____, _____, and _____.
- _____

Numerical Response

2. To the nearest tenth, the energy released when 1.00 mol of AgI_(s) is formed from its elements is _____ kJ.

Use the following information to answer the next two questions.

Glucose is a biological fuel used by cells to satisfy the energy needs of plants and animals. The overall reaction for the metabolism of glucose is represented by the **unbalanced** equation



4. The balanced equation and the enthalpy change for the cellular respiration of glucose can be represented as

- A. $\text{C}_6\text{H}_{12}\text{O}_6(\text{s}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l}) + 593.8 \text{ kJ}$
B. $\text{C}_6\text{H}_{12}\text{O}_6(\text{s}) + 6 \text{ O}_2(\text{g}) + 2802.7 \text{ kJ} \rightarrow 6 \text{ CO}_2(\text{g}) + 6 \text{ H}_2\text{O}(\text{l})$
C. $\text{C}_6\text{H}_{12}\text{O}_6(\text{s}) + 6 \text{ O}_2(\text{g}) \rightarrow 6 \text{ CO}_2(\text{g}) + 6 \text{ H}_2\text{O}(\text{l}) + 2802.7 \text{ kJ}$
D. $\text{C}_6\text{H}_{12}\text{O}_6(\text{s}) + 6 \text{ O}_2(\text{g}) \rightarrow 6 \text{ CO}_2(\text{g}) + 6 \text{ H}_2\text{O}(\text{l}) + 2538.7 \text{ kJ}$

5. If solid glucose is completely burned in the flame of a Bunsen burner, the enthalpy change is

- A. greater than it is during cellular respiration because the production of H₂O_(g) releases more energy than does the production of H₂O_(l)
B. less than it is during cellular respiration because the production of H₂O_(g) releases less energy than does the production of H₂O_(l)
C. the same as it is in the body because the enthalpy change is independent of the state of the products
D. the same as it is in cellular respiration because they are identical processes


6. When 1.65 g of ethanal (CH₃CHO_(l)) is burned in a calorimeter to produce H₂O_(l) and CO₂_(g), 44.7 kJ of heat energy is produced. According to this experimental data, the molar enthalpy of combustion of ethanal is

- A. $+1.52 \times 10^3 \text{ kJ/mol}$ B. -76.6 kJ/mol C. -165 kJ/mol D. $-1.19 \times 10^3 \text{ kJ/mol}$

Numerical Response

3. To the nearest hundredth, A student heated a 120.0 g sample of H₂O_(l) from 21.0°C to 32.5°C by adding 5.93 kJ of energy. The student then used this data to calculate the specific heat capacity of water and compared it with the standard value. The experimental percentage difference was _____ %.

Use the following information to answer the next question.




**Numerical Response**

4. To the nearest hundredth, the amount of energy released by the combustion of 100 g of C₂H₂_(g) is _____ MJ.


Use the following information to answer the next question.

Many insects and small animals have unique defence systems. Bombardier beetles fight off predators with a hot chemical spray. This spray consists of solutions of hydroquinone ($\text{C}_6\text{H}_4(\text{OH})_2(aq)$), hydrogen peroxide ($\text{H}_2\text{O}_2(aq)$), and enzymes, which are secreted by the beetles' glands.

Reaction Equation Related to Spray Formation






A chemical reaction that occurs in order to produce the hot chemical spray can be represented by the equation



7. The heat of reaction for the production of this hot chemical spray is
- A. -489.2 kJ B. -203.4 kJ C. -82.4 kJ D. $+12.2 \text{ kJ}$


Use the following equations to answer the next question.

Energy Reaction Equations



8. The energy reactions above involve the conversion of energy for metabolic (body) processes. The chronological order of these reactions is
- A. I, III, and II B. III, II, and I C. II, III, and I D. II, I, and III
9. The total enthalpy change associated with the conversion of 1.00 Mg of water at 20.0°C into steam at 250.0°C could be calculated by using the formula
- A. $[1.00 \text{ Mg} \times 4.19 \text{ J}/(\text{g} \cdot ^\circ\text{C}) \times 80.0^\circ\text{C}] + [(1.00 \text{ Mg}/18.02 \text{ g/mol}) \times 40.65 \text{ kJ/mol}]$
B. $[1.00 \text{ Mg} \times 2.02 \text{ J}/(\text{g} \cdot ^\circ\text{C}) \times 230.0^\circ\text{C}] + [(1.00 \text{ Mg}/18.02 \text{ g/mol}) \times 40.65 \text{ kJ/mol}]$
C. $[1.00 \text{ Mg} \times 4.19 \text{ J}/(\text{g} \cdot ^\circ\text{C}) \times 80.0^\circ\text{C}] + [(1.00 \text{ Mg}/18.02 \text{ g/mol}) \times 40.65 \text{ kJ/mol}]$
 $+ [1.00 \text{ Mg} \times 4.19 \text{ J}/(\text{g} \cdot ^\circ\text{C}) \times 150.0^\circ\text{C}]$
D. $[1.00 \text{ Mg} \times 4.19 \text{ J}/(\text{g} \cdot ^\circ\text{C}) \times 80.0^\circ\text{C}] + [(1.00 \text{ Mg}/18.02 \text{ g/mol}) \times 40.65 \text{ kJ/mol}]$
 $+ [1.00 \text{ Mg} \times 2.02 \text{ J}/(\text{g} \cdot ^\circ\text{C}) \times 150.0^\circ\text{C}]$

At the Acme Gas Plant in Texas, environmental and economic concerns have resulted in the development of an efficient process for the removal of sulfur from sour gas, which is a mixture of hydrocarbons and $\text{H}_2\text{S}_{(g)}$. In the first step of the process, one-third of the $\text{H}_2\text{S}_{(g)}$ reacts with $\text{O}_{2(g)}$ to produce $\text{SO}_{2(g)}$. In the second step of the process, the $\text{SO}_{2(g)}$ produced reacts with the remaining $\text{H}_2\text{S}_{(g)}$ to form elemental sulfur and water.




To maximize the amount of sulfur removed from the sour gas, the gas plant engineers apply Le Châtelier's principle.

10. According to the overall equilibrium equation above, the amount of sulfur removed may be increased by
- | | |
|--|---|
| A. adding a catalyst | B. removing water vapour |
| C. increasing the volume of the system | D. increasing the temperature of the system |
11. As $\text{H}_2\text{S}_{(g)}$ forms $\text{S}_{(s)}$, the oxidation number of sulfur
- changes from 0 to -2 and sulfur is reduced
 - changes from -2 to 0 and sulfur is oxidized
 - decreases by 2 and hydrogen sulfide acts as the reducing agent
 - stays the same because the sulfur is neither oxidized nor reduced


Use the following information to answer the next question.

The sulfur produced in step II is initially produced in liquid form. As it cools, it is converted from a liquid state to a solid state as represented by the equation



12. In terms of energy, this conversion is
- | | |
|--|---|
| A. endothermic, releases heat, and has a positive ΔH | B. exothermic, releases heat, and has a negative ΔH |
| C. exothermic, absorbs heat, and has a negative ΔH | D. endothermic, absorbs heat, and has a positive ΔH |

At one time, an aqueous solution of formaldehyde called formalin ($\text{CH}_2\text{O}_{(aq)}$) was used as a disinfectant and as a tissue preservative. Today, formalin is commonly used in the industrial preparation of plastics and resins. Formalin can be produced by reacting methanol with acidified potassium dichromate, as represented by the following **unbalanced** equation.



13. The type of reaction that this equation represents is
- | | |
|--|------------------------------------|
| A. a Brønsted–Lowry acid–base reaction | B. an oxidation–reduction reaction |
| C. a formation reaction | D. a combustion reaction |

14. When the above equation is balanced, the equation is

- A. $\text{CH}_3\text{OH}_{(l)} + \text{Cr}_2\text{O}_7^{2-}{}_{(aq)} + 14 \text{H}^+{}_{(aq)} \rightarrow \text{CH}_2\text{O}_{(aq)} + 2 \text{Cr}^{3+}{}_{(aq)} + 7 \text{H}_2\text{O}_{(l)}$
B. $3 \text{CH}_3\text{OH}_{(l)} + \text{Cr}_2\text{O}_7^{2-}{}_{(aq)} + 14 \text{H}^+{}_{(aq)} \rightarrow 3 \text{CH}_2\text{O}_{(aq)} + 2 \text{Cr}^{3+}{}_{(aq)} + 7 \text{H}_2\text{O}_{(l)}$
C. $3 \text{CH}_3\text{OH}_{(l)} + \text{Cr}_2\text{O}_7^{2-}{}_{(aq)} + 8 \text{H}^+{}_{(aq)} \rightarrow 3 \text{CH}_2\text{O}_{(aq)} + 2 \text{Cr}^{3+}{}_{(aq)} + 7 \text{H}_2\text{O}_{(l)}$
D. $3 \text{CH}_3\text{OH}_{(l)} + \text{Cr}_2\text{O}_7^{2-}{}_{(aq)} + 8 \text{H}^+{}_{(aq)} \rightarrow 3 \text{CH}_2\text{O}_{(aq)} + 2 \text{Cr}^{3+}{}_{(aq)} + 8 \text{H}_2\text{O}_{(l)}$

Use your recorded answer for **Multiple Choice 14** to answer **Numerical Response 5**.*

Numerical Response

5. To the nearest tenth, when 39.5 kg of methanol is reacted, the mass of formalin produced is _____ kg.

*You can receive marks for this question even if the previous question was answered incorrectly.

15. Iodine solutions, which contain a suspension of $\text{I}_2_{(s)}$ have a brown colour. Which of the following metals will **not** cause an iodine solution to change colour?

- A. $\text{Ni}_{(s)}$ B. $\text{Cu}_{(s)}$ C. $\text{Ag}_{(s)}$ D. $\text{Mg}_{(s)}$

Use the following information to answer the next question.

A sample of $\text{Na}_2\text{S}_2\text{O}_3_{(aq)}$ is titrated with acidified $\text{KMnO}_4_{(aq)}$ a pink endpoint. One product of this redox reaction is $\text{SO}_4^{2-}{}_{(aq)}$.

16. A product of the reduction half-reaction is

- A. $\text{H}^+{}_{(aq)}$ B. $\text{Mn}^{2+}{}_{(aq)}$ C. $\text{SO}_4^{2-}{}_{(aq)}$ D. $\text{S}_2\text{O}_3^{2-}{}_{(aq)}$

Use the following information to answer the next question.

In an experiment, dilute hydrochloric acid was added to a test tube containing several small pieces of $\text{Zn}_{(s)}$.

17. Which of the following observations could have been made during this experiment?

- A. There was no reaction. B. A thick white precipitate formed.
C. A colourless gas was produced and the test tube cooled off.
D. A colourless gas was produced and the test tube warmed up.

Use the following information to answer the next question.





A student used an acidified 6.31×10^{-2} mol/L $\text{KMnO}_4_{(aq)}$ solution to titrate 25.0 mL samples of $\text{Fe}^{2+}{}_{(aq)}$ solution of unknown concentration. In the reactions, the $\text{Fe}^{2+}{}_{(aq)}$ ion was oxidized to the $\text{Fe}^{3+}{}_{(aq)}$ ion. The student completed five trials and summarized the data in the table.

Trial Number	1	2	3	4	5
Final Buret Reading (mL)	17.55	35.65	26.40	42.65	16.85
Initial Buret Reading (mL)	0.30	17.55	10.05	26.40	0.55
Final Colour	purple	purple	pink	pink	pink

18. According to the student's data, the concentration of $\text{Fe}^{2+}{}_{(aq)}$


- A. 0.206 mol/L B. 0.218 mol/L C. 0.213 mol/L D. 0.223 mol/L

19. The half-reaction to which all other half-cell potentials are compared is



Use the following information to answer the next two questions.

Electronic hobbyists often “etch” circuit boards. In this process, unwanted copper foil is removed from a copper-clad plastic circuit board by immersing the board in a bath of iron (III) chloride solution. The equation for the net reaction is



20. In the reaction above, in which the unwanted copper foil is removed,

A. copper ions are reduced

B. copper atoms are oxidized

C. iron (II) ions act as the oxidizing agent

D. iron (III) ions act as the reducing agent

21. Which of the following statements and corresponding net voltages are correct for this reaction?

A. It is a spontaneous reaction with an $E^\circ_{\text{net}} = +0.43\text{ V}$.


B. It is a spontaneous reaction with an $E^\circ_{\text{net}} = +1.11\text{ V}$.

C. A power supply is required because the $E^\circ_{\text{net}} = -0.43\text{ V}$.

D. A power supply is required because the $E^\circ_{\text{net}} = -1.11\text{ V}$.

Use the following information to answer the next two questions.

Hydrogen–oxygen fuel cells have been used for years in spacecraft and more recently in small-scale power plants to generate electricity. Now, some governments and companies are working together to perfect this type of fuel cell for automobile use, and experiments are currently being conducted with operational prototypes. A diagram of a hydrogen–oxygen fuel cell is shown below.




Numerical Response

6. In the diagram above, the anode, the cathode, the electrolyte, and a product of the reaction are labeled, respectively, _____, _____, _____, and _____.

22. From an ecological perspective, a reason why hydrogen–oxygen fuel cells should **not** be used to power automobiles is that
- A. hydrogen fuel can be produced through the electrolysis of seawater by using the energy produced from burning fossil fuels
 - B. cars powered by a hydrogen–oxygen fuel cell would be up to 30% more efficient than cars powered by gasoline
 - C. water vapour is the primary byproduct of the cell
 - D. oxygen is readily available from the atmosphere
-

Use the following information to answer the next two questions.

Copper can be refined (purified) using an apparatus like the one shown below, which is a small-scale version of an industrial apparatus.




23. In this electrochemical cell, the purified copper sheet acts as the
- A. anode and is the site where $\text{SO}_4^{2-}_{(aq)}$ ions are oxidized
 - B. cathode and is the site where $\text{SO}_4^{2-}_{(aq)}$ ions are reduced
 - C. anode and is the site where $\text{Cu}^{2+}_{(aq)}$ ions are oxidized
 - D. cathode and is the site where $\text{Cu}^{2+}_{(aq)}$ ions are reduced

Numerical Response


7. If the direct current power supply produces a steady 3.50 A current, then to the nearest tenth, the time required to deposit 0.100 g of purified copper is _____ s.
-

Use the following information to answer the next three questions.

The silver oxide alkaline cell is a miniature power source used in watches, calculators, hearing aids, and cameras. The construction of this cell is shown in the following diagram.



Half-Reactions




24. During the discharging of the cell, the substance oxidized is
- A. $\text{Zn}(s)$ B. $\text{Ag}(s)$ C. $\text{H}_2\text{O}(l)$ D. $\text{Ag}_2\text{O}(s)$
25. In this cell, the separator must be porous in order to
- A. allow migration of ions B. replenish the electrolyte
C. provide a pathway for electron flow D. provide a surface on which electron transfer can occur

Numerical Response

8. During discharge, the voltage generated by the cell, to the nearest hundredth, is +/- _____ V.

Use the following information to answer the next question.



26. The cell in the diagram was constructed and connected by a chemistry student. The voltage of the cell remained at 0.00 V trial after trial. One possible reason for the malfunction of the cell was that the
- A. concentrations of the solutions were too low B. solution in the U-tube was a non-electrolyte
 C. redox reaction was non-spontaneous D. voltmeter was connected backward

Use the following information to answer the next question.

A Midwest company processes aluminum “logs” for commercial use. The first step in the process involves removing the natural aluminum oxide coating from the logs.


27. Once the protective coating has been removed, the $\text{Al}_{(s)}$ surface undergoes a redox reaction with $\text{H}_2\text{O}_{(l)}$. In this reaction,
- A. $\text{H}_2(g)$ is evolved and the solution becomes basic B. $\text{O}_2(g)$ is evolved and the solution becomes basic
 C. $\text{H}_2(g)$ is evolved and the solution becomes acidic D. $\text{O}_2(g)$ is evolved and the solution becomes acidic

Use the following information to answer the next question.

In the late 1980s, the Canadian dollar bill was replaced by a coin commonly called the “loonie.” The loonie is manufactured from nickel disks that are stamped and then coated with a thin layer of copper (87.5%) and tin (12.5%) to provide the shiny gold-coloured appearance. This layer is applied through an electrolysis process in which the stamped loonie is one of the electrodes and copper metal is the other electrode.

28. If the plating of the loonie occurs in a $\text{Sn}^{2+}_{(aq)}$ and $\text{Cu}^{2+}_{(aq)}$ solution, then the reaction that occurs at the cathode is
- A. $2 \text{H}_2\text{O}_{(l)} + 2e^- \rightarrow \text{H}_2(g) + 2 \text{OH}^-_{(aq)}$ B. $2 \text{H}_2\text{O}_{(l)} \rightarrow \text{O}_2(g) + 4 \text{H}^+_{(aq)} + 4e^-$
 C. $\text{Cu}^{2+}_{(aq)} + 2e^- \rightarrow \text{Cu}_{(s)}$ D. $\text{Cu}_{(s)} \rightarrow \text{Cu}^{2+}_{(aq)} + 2e^-$

29. Of the graphs below, the one that best illustrates the relationship between $[\text{H}_3\text{O}^+_{(aq)}]$ and $[\text{OH}^-_{(aq)}]$ in a solution is



30. If equal moles of acid and base are mixed, then which of the following pairs of species yields a solution with a pH closest to that of pure water at 25°C ?
- A. $\text{HSO}_4^-_{(aq)}$ and $\text{OH}^-_{(aq)}$ B. $\text{H}_2\text{S}_{(aq)}$ and $\text{OH}^-_{(aq)}$
 C. $\text{H}_3\text{O}^+_{(aq)}$ and $\text{HCO}_3^-_{(aq)}$ D. $\text{H}_3\text{O}^+_{(aq)}$ and $\text{OH}^-_{(aq)}$

Use the following information to answer the next question.


The labels came off four cleaning solution containers found under a kitchen sink. Each of the cleaning solutions was tested with two available indicators, and the following results were recorded.

Cleaning Solution	Bromothymol Blue	Phenolphthalein
1	blue	pink
2	blue	colourless
3	green	colourless
4	blue	light pink

Numerical Response

9. Listed in order from lowest to highest pH, the cleaning solutions are, respectively, _____, _____, _____, and _____.

CleanChem Inc. in Nevada is a bulk manufacturer of concentrated bleach ($\text{NaOCl}_{(aq)}$). The bleach reacts with water to form a solution with a pH of 10.87.



31. In this reaction, the substances that act as Brønsted–Lowry acids are
A. $\text{OCl}^{-}_{(aq)}$ and $\text{H}_2\text{O}_{(l)}$ B. $\text{OCl}^{-}_{(aq)}$ and $\text{HOCl}_{(aq)}$ C. $\text{OCl}^{-}_{(aq)}$ and $\text{OH}^{-}_{(aq)}$ D. $\text{H}_2\text{O}_{(l)}$ and $\text{HOCl}_{(aq)}$
32. The substance in the equation above that may act as an amphoteric species is
A. $\text{OCl}^{-}_{(aq)}$ B. $\text{H}_2\text{O}_{(l)}$ C. $\text{HOCl}_{(aq)}$ D. $\text{OH}^{-}_{(aq)}$
33. The two species in equimolar amounts that could act as a buffer in this bleach solution are
A. $\text{OCl}^{-}_{(aq)}$ and $\text{HOCl}_{(aq)}$ B. $\text{HOCl}_{(aq)}$ and $\text{OH}^{-}_{(aq)}$ C. $\text{OCl}^{-}_{(aq)}$ and $\text{H}_2\text{O}_{(l)}$ D. $\text{H}_2\text{O}_{(l)}$ and $\text{OH}^{-}_{(aq)}$
34. In this bleach solution, the acid–base indicator
A. phenolphthalein would be colourless B. alizarin yellow R would be orange
C. indigo carmine would be green D. methyl orange would be red

Use the following information to answer the next question.

A bleach solution can be made by dissolving chlorine gas in a sodium hydroxide solution, as shown by the equation



35. Mixing a bleach solution with an acid solution can be dangerous because it can cause
A. an increase in pH in the bleach solution
B. a shift in the equilibrium to the products
C. an increase in $\text{Cl}_2_{(g)}$ concentration in the bleach solution
D. an increase in $\text{Cl}^{-}_{(aq)}$ concentration in the bleach solution

Use the following information to answer the next two questions.

Coal is composed of many organic substances. When coal is mixed with water, acids are formed from the impurities found in the coal. Technicians at Southampton Coal Corporation refer to this mixture of acids as “humic acid.”

36. A standard quality-control test involves the titration of monoprotic humic acid with $\text{NaOH}_{(aq)}$. If a 10.0 mL sample of a saturated solution of humic acid reacts with 15.9 mL of a 0.100 mol/L $\text{NaOH}_{(aq)}$ solution, then the concentration of the acid is
- A. 0.0629 mol/L B. 0.100 mol/L C. 0.159 mol/L D. 0.059 mol/L

Use your recorded answer for **Multiple Choice 36** to answer **Multiple Choice 37**.*

37. If the pH of a specific humic acid sample is 4.50, then the K_a value will be
- A. 1.0×10^{-8} B. 1.6×10^{-8} C. 1.7×10^{-8} D. 6.3×10^{-9}

*You can receive marks for this question even if the previous question was answered incorrectly.

38. Chloroacetic acid ($\text{CH}_2\text{ClCOOH}_{(aq)}$) has a $K_a = 1.4 \times 10^{-3}$. This acid could best be described as a
- A. weak inorganic acid B. diprotic organic acid
C. weak monoprotic acid D. strong monoprotic acid
39. In the equation $\text{HNO}_3_{(aq)} + \text{N}_2\text{H}_4_{(aq)} \rightleftharpoons \text{NO}_3^-_{(aq)} + \text{N}_2\text{H}_5^+_{(aq)}$, one conjugate acid–base pair is
- A. $\text{HNO}_3_{(aq)}$ and $\text{N}_2\text{H}_5^+_{(aq)}$ B. $\text{HNO}_3_{(aq)}$ and $\text{N}_2\text{H}_4_{(aq)}$
C. $\text{N}_2\text{H}_4_{(aq)}$ and $\text{N}_2\text{H}_5^+_{(aq)}$ D. $\text{N}_2\text{H}_4_{(aq)}$ and $\text{NO}_3^-_{(aq)}$

Use the following information to answer the next question.





A 0.500 mol/L solution of hydrazine ($\text{N}_2\text{H}_4_{(aq)}$) contains the following equilibrium concentrations.
 $[\text{N}_2\text{H}_4_{(aq)}] = 0.498 \text{ mol/L}$ $[\text{OH}^-_{(aq)}] = 2.14 \times 10^{-3} \text{ mol/L}$ $[\text{N}_2\text{H}_5^+_{(aq)}] = 2.14 \times 10^{-3} \text{ mol/L}$

Numerical Response

10. The K_b for hydrazine, in scientific notation, is $a.bc \times 10^{-d}$. The values of a , b , c , and d are, respectively, _____, _____, _____, and _____.

Use the following information to answer the next question.

When equally concentrated solutions of $\text{HNO}_3(aq)$, $\text{CH}_3\text{COOH}(aq)$, $\text{HOOC}\text{COOH}(aq)$, and $\text{Na}_2\text{S}(aq)$ were titrated with either a strong acid or strong base, the following titration curves were obtained.




Numerical Response

11. Match each of the graphs, as numbered above, with the corresponding titration species listed below.

$\text{HNO}_3(aq)$	_____	(Record as the first digit)
$\text{CH}_3\text{COOH}(aq)$	_____	(Record as the second digit)
$\text{HOOC}\text{COOH}(aq)$	_____	(Record as the third digit)
$\text{Na}_2\text{S}(aq)$	_____	(Record as the fourth digit)

Use the following information to answer the next question.

The burning of methane in a Bunsen burner to produce energy can be represented by the equation



40. A student determined that the reaction represented by the equation above is **not** at equilibrium because

- | | |
|---------------------------------------|---|
| A. the system is open | B. a catalyst is not present |
| C. the temperature is constant | D. both reactants and products are gases |

45. The electron configuration of a ground-state copper atom is
 A. $[\text{Ar}]4s^24d^4$. B. $[\text{Ar}]4s^23d^9$. C. $[\text{Ar}]3d^9$. D. $[\text{Ar}]4s^13d^{10}$.
46. Which of these choices is the electron configuration of an *excited state* of an iron atom?
 A. $[\text{Ar}]4s^23d^6$ B. $[\text{Ar}]3d^5$ C. $[\text{Ar}]4s^13d^7$ D. $[\text{Ar}]4s^13d^5$
47. Which of these pairs consists of *isoelectronic* species?
 A. Zn^{2+} and Cu^{2+} B. Na^+ and K^+ C. Cl^- and S D. K^+ and Cl^-
48. The electron configuration of a cobalt (III) ion is
 A. $[\text{Ar}]3d^5$. B. $[\text{Ar}]4s^13d^5$. C. $[\text{Ar}]4s^23d^4$. D. $[\text{Ar}]3d^6$.
49. Arrange these ions in order of increasing ionic radius: K^+ , P^{3-} , S^{2-} , Cl^- .
- Increasing radius \rightarrow
- Row 1 $\text{K}^+ < \text{Cl}^- < \text{S}^{2-} < \text{P}^{3-}$
 Row 2 $\text{K}^+ < \text{P}^{3-} < \text{S}^{2-} < \text{Cl}^-$
 Row 3 $\text{P}^{3-} < \text{S}^{2-} < \text{Cl}^- < \text{K}^+$
 Row 4 $\text{Cl}^- < \text{S}^{2-} < \text{P}^{3-} < \text{K}^+$
- A. Row 1 B. Row 2 C. Row 3 D. Row 4
50. Which of these elements has the smallest ionization energy?
 A. Li B. Na C. K D. Rb
51. Which of these elements has the greatest electron affinity (largest positive value)?
 A. Al B. Si C. P D. S
52. The total number of bonding electrons in a molecule of formaldehyde (H_2CO) is
 A. 3. B. 4. C. 6. D. 8.
53. Which molecule has a Lewis structure that does not obey the octet rule?
 A. CS_2 B. NO_2 C. PH_3 D. CCl_4
54. Which of the following substances should have the highest boiling point?
 A. CH_4 B. Cl_2 C. CH_3Cl D. Kr
55. Which two properties are more typical of molecular compounds than of ionic compounds?
 I. They are gases or liquids at room temperature. II. They have high melting points.
 III. Solids do not conduct electricity, but liquids do. IV. Atoms share electrons.
- A. I and IV B. I and III C. II and III D. II and IV
56. Which one of the following substances should exhibit hydrogen bonding in the liquid state?
 A. PH_3 B. H_2S C. CH_4 D. NH_3

57. A student was given data concerning the boiling points of hydrogen compounds in the fourth period of the periodic table.

Compound	Boiling Point (°C)
GeH ₄	-89
AsH ₃	-55
H ₂ Se	-42
HBr	-67

The best hypothesis the student could make to explain the drop in boiling points between H₂Se and HBr is that

- A. the H₂Se intramolecular bonds are more polar than for HBr
- B. hydrogen bonding occurs with H₂Se but not with HBr
- C. fewer intermolecular bonds can form with HBr compared to H₂Se because of its shape
- D. HBr has too many lone pairs of electrons to make strong intermolecular bonds

Part B: Written Response


Use the following information to answer the next question.

Most arenas and curling rink have artificial ice. Many ice-making plants use ammonia as the refrigerant. The ammonia is circulated in pipes under the ice of the arena or curling rink. For this question, assume that the **only** changes to the ammonia are represented in the equilibrium.




1. a. On the axes provided, draw and label, as precisely as possible, the graphs that represent the energy changes that occur to the ammonia below the ice surface and to the water on the ice surface as the refrigeration system operates. Assume that the water applied to the ice surface is initially at 20.00°C.

Ammonia Change



Water Change



- b. What mass of ammonia must undergo a phase change in order to change 1.00×10^7 g (10.0 kL) of water at 20.00°C to ice at 0.00°C?

Use the following information to answer the next question.

The formation of a pollutant gas, nitrogen monoxide ($\text{NO}_{(g)}$), by the reaction of nitrogen with oxygen in a gasoline engine can be affected by changing the combustion temperature within the engine. The equilibrium constant for the production of one mole of $\text{NO}_{(g)}$ at 25°C is 1.0×10^{-17}

2. Explain how an increase in temperature could affect the concentration of the pollutant gas and the equilibrium constant.

Your response should also include

- relevant chemical equation(s) and values from the chemistry data booklet
 - a description of two ways that car manufacturers could reduce the $\text{NO}_{(g)}$ emissions in new model vehicles
3. For the following compounds or ions, draw the Lewis dot diagram and predict its molecular geometry. Indicate any compound that has resonance structures.

a. PF_5

b. SiCl_4

c. NO_3^-

Answers

Multiple Choice

1. B 2. B 3. A 4. C 5. B 6. D 7. B 8. C 9. D 10. B 11. B
 12. B 13. B 14. C 15. C 16. B 17. D 18. A 19. D 20. B 21. A 22. A
 23. D 24. A 25. A 26. B 27. A 28. C 29. A 30. D 31. D 32. B 33. A
 34. B 35. C 36. C 37. D* 38. C 39. C 40. A 41. D 42. D 43. D[†] 44. C
 45. D 46. C 47. D 48. D 49. C 50. D 51. D 52. D 53. B 54. C 55. A
 56. D 57. C

Numerical Response

1. 3214 2. 61.8 3. 2.56 or 2.63 4. 4.82 5. 37.0 6. 2654
 7. 86.8 8. 1.59 9. 3241 10. 9206 11. 2413 12. 2.54[§]

Linked Items:


*If MC36 is A, then MC37 is B
 If MC36 is B, then MC37 is A
 If MC36 is C, then MC37 is D*
 If MC36 is D, then MC37 is C

[†]If MC42 is A, then MC37 is A
 If MC42 is B, then MC37 is D
 If MC42 is C, then MC37 is A
 If MC42 is D, then MC37 is D[†]

[§]If MC43 is A, then NR12 is 5.09
 If MC43 is B, then NR12 is 4.62
 If MC43 is C, then NR12 is 2.31
 If MC43 is D, then NR12 is 2.54[§]

Written Response

1. a.



b. Heat Gained (Ammonia) = Heat Lost (Water)

$$n_{\text{NH}_3} \Delta H_{\text{vap, NH}_3} = m_w C_w \Delta T_w + n_w \Delta H_{\text{fus}}$$


$$n_{\text{NH}_3} = \frac{m_w C_w \Delta T_w + n_w \Delta H_{\text{fus}}}{\Delta H_{\text{vap, NH}_3}} = \frac{(1.00 \times 10^4 \text{ kg})(4.19 \text{ kJ}/(\text{kg} \cdot ^\circ \text{C}))(20.00^\circ \text{C}) + \left(\frac{1.00 \times 10^7 \text{ g}}{18.02 \text{ g/mol}}\right)(6.03 \text{ kJ/mol})}{(23.3 \text{ kJ/mol})}$$

$$n_{\text{NH}_3} = 179582.9145 \text{ mol}$$

$$m_{\text{NH}_3} = nM = (179582.9145 \text{ mol})(17.04 \text{ g/mol}) = 3,060,092.864 \text{ g}$$

$$m_{\text{NH}_3} = 3.06 \times 10^6 \text{ g} \quad \text{or} \quad 3.06 \times 10^3 \text{ kg} \quad \text{or} \quad 3.06 \text{ Mg}$$

2. Chemical Equation



Explanations

The reaction is endothermic. Therefore, according to Le Châtelier's Principle, increasing the temperature should shift the equilibrium to the right that will cause an increase in $[\text{NO}(\text{g})]$.


The K_{eq} value is temperature dependent. At a higher temperature, more $\text{NO}(\text{g})$ is produced therefore the K value would increase.

$$K_{\text{eq}} = \frac{[\text{NO}(\text{g})]^\uparrow}{[\text{N}_2(\text{g})]^\downarrow \times [\text{O}_2(\text{g})]^\downarrow} = 1.0 \times 10^{-17} \text{ (increases)}$$

Ways to reduce $\text{NO}(\text{g})$ emissions


- Car manufacturers should try to decrease engine combustion temperatures.
- Install emission control devices. (Catalytic reduction of $\text{NO}(\text{g})$ emissions.)
- Install $\text{N}_2(\text{g})$ absorbent before combustion takes place in the engine.
- More efficient fuel/car with an explanation.
- Use of hybrid/electric/solar cars.

3. a. PF_5




Trigonal bipyramid

b. SiCl_4



Tetrahedral

c. NO_3^-



Trigonal planar